

Boundary Value Problem for $r^2 d^2 f/dr^2 + f = f^3$ (III): Global Solution and Asymptotics

Chie Bing Wang*

*Department of Mathematics, University of Pittsburgh
 Pittsburgh, PA 15260.*

Abstract

Based on the results in the previous papers that the boundary value problem $y'' - y' + y = y^3$, $y(0) = 0$, $y(\infty) = 1$ with the condition $y(x) > 0$ for $0 < x < \infty$ has a unique solution $y^*(x)$, and $a^* = y^{*\prime}(0)$ satisfies $0 < a^* < 1/4$, in this paper we show that $y'' - y' + y = y^3$, $-\infty < x < 0$, with the initial conditions $y(0) = 0$, $y'(0) = a^*$ has a unique solution by using functional analysis method. So we get a globally well defined bounded function $y^*(x)$, $-\infty < x < +\infty$. The asymptotics of $y^*(x)$ as $x \rightarrow -\infty$ and as $x \rightarrow +\infty$ are obtained, and the connection formulas for the parameters in the asymptotics and the numerical simulations are also given. Then by the properties of $y^*(x)$, the solution to the boundary value problem $r^2 f'' + f = f^3$, $f(0) = 0$, $f(\infty) = 1$ is well described by the asymptotics and the connection formulas.

1 Introduction

In this paper, we use the results obtained in [2] [3] to study the boundary value problem

$$r^2 f'' + f = f^3, \quad 0 < r < \infty, \quad (1.1)$$

$$f(r) \rightarrow 0, \text{ as } r \rightarrow 0, \quad (1.2)$$

$$f(\infty) = 1. \quad (1.3)$$

* Current address: Department of Mathematics, University of California, Davis, CA 95616. e-mail: cbwang@math.ucdavis.edu

If we make a transformation $r = e^x$, $f(r) = y(x)$, the equation is changed to $y'' - y' + y = y^3$.

The results in [2] [3] can be briefly summarized as follows. The following boundary value problem

$$\begin{aligned} & y'' - y' + y = y^3, \quad 0 < x < \infty, \\ (P^+) \quad & y(0) = 0, y(\infty) = 1, \\ & y(x) > 0, \quad 0 < x < \infty, \end{aligned}$$

has a unique solution $y^*(x)$ for $0 < x < \infty$. The formula of

$$a^* = y^{*'}(0), \tag{1.4}$$

is obtained, and

$$0 < a^* < 1/4. \tag{1.5}$$

In this paper, we show that the initial value problem

$$\begin{aligned} & y'' - y' + y = y^3, \quad -\infty < x < 0, \\ (P^-) \quad & y(0) = 0, \\ & y'(0) = a^*, \end{aligned}$$

has a unique solution by discussing the corresponding integral equation on a Banach space. Then we get a global solution $y^*(x)$ of $y'' - y' + y = y^3$ for $-\infty < x < +\infty$ and the asymptotics as $x \rightarrow -\infty$ and as $x \rightarrow +\infty$. And any bounded solution $y(x)$ of this equation satisfying $y(-\infty) = 0$, $y(+\infty) = 1$ is expressed as $y(x) = y^*(x - \tau)$, where τ is the largest zero of $y(x)$. Therefore we have solved the boundary value problem (1.1) (1.2) (1.3).

This paper is organized as follows. In Sect. 2, $y^*(x)$ is extended to the negative axis by using contraction mapping theorem. In Sect. 3 we give numerical approximations for some important values including a^* , which are used to represent asymptotics of the solution. In Sect. 4, the asymptotic expressions and connection formulas are given.

2 Global Solution

In [2], we have proved problem (P^+) has a unique solution $y^*(x) = y(x, a^*)$. In this section we want to extend the solution to the negative axis. Consider the following problem

$$y'' - y' + y = y^3, \quad -\infty < x < 0, \tag{2.1}$$

$$(P^-) \quad y(0) = 0, \tag{2.2}$$

$$y'(0) = a^*. \tag{2.3}$$

Let

$$t = -x, \quad u(t) = e^{-x/2}y(x).$$

Then problem (P^-) is equivalent to the following integral equation

$$u(t) = -\frac{2}{\sqrt{3}}a^* \sin \frac{\sqrt{3}}{2}t + \frac{2}{\sqrt{3}} \int_0^t e^{-s} \sin \frac{\sqrt{3}}{2}(t-s)u^3(s) ds, \quad (2.4)$$

where $t \geq 0$. Define

$$X = \{f(t) | f \text{ is continuous on } [0, \infty), |f| \leq \frac{1}{2}, t \in [0, \infty)\}, \quad (2.5)$$

$$d(f, g) = \sup_{t \in [0, \infty)} |f(t) - g(t)|, \quad (2.6)$$

for $f, g \in X$. It is not difficult to prove the following lemma.

Lemma 1 (X, d) is a complete metric space.

Define

$$T(u)(t) = -\frac{2}{\sqrt{3}}a^* \sin \frac{\sqrt{3}}{2}t + \frac{2}{\sqrt{3}} \int_0^t e^{-s} \sin \frac{\sqrt{3}}{2}(t-s)u^3(s) ds, \quad (2.7)$$

for $u \in X$.

Theorem 1 T has precisely one fixed point u^* in X .

Proof. Let us first prove $T(u) \in X$ when $u \in X$. By (2.7), $T(u)(t)$ is continuous on $[0, \infty)$. By (1.5),

$$a^* < \frac{1}{4},$$

which implies that if $|u| \leq 1/2$,

$$|T(u)(t)| \leq \frac{2}{\sqrt{3}}a^* + \frac{2}{\sqrt{3}} \left(\frac{1}{2}\right)^3 < \frac{\sqrt{3}}{4} < \frac{1}{2}.$$

So $T(u) \in X$.

Next we show T is a contraction on X . In fact, if $u_1, u_2 \in X$, there is

$$\begin{aligned} |T(u_1) - T(u_2)| &= \left| \frac{2}{\sqrt{3}} \int_0^t e^{-s} \sin \frac{\sqrt{3}}{2}(t-s)(u_1 - u_2)(u_1^2 + u_1u_2 + u_2^2) ds \right| \\ &\leq \frac{3}{2\sqrt{3}} \sup_{t \in [0, \infty)} |u_1(t) - u_2(t)| \\ &\leq \frac{\sqrt{3}}{2} d(u_1, u_2), \end{aligned}$$

which implies

$$d(T(u_1), T(u_2)) \leq \frac{\sqrt{3}}{2} d(u_1, u_2).$$

So T is a contraction. By Banach fixed point theorem, this theorem is proved. $\square$

Theorem 2 *There is a unique solution y^* to the problem*

$$y'' - y' + y = y^3, \quad -\infty < x < \infty, \quad (2.8)$$

$$(P) \quad y(0) = 0, \quad y(\infty) = 1, \quad (2.9)$$

$$y(x) > 0, \quad 0 < x < \infty. \quad (2.10)$$

And y^* has infinitely many zeros $x_n (n = 0, 1, 2, \dots)$

$$-\infty < \dots < x_n < \dots < x_1 < x_0 = 0,$$

also

$$y(-\infty) = 0, \quad |y^*(x)| \leq \frac{1}{2},$$

for $x \in (-\infty, 0]$.

Proof. The Theorem 2 in [2] and Theorem 1 imply that problem (P) has a unique solution y^* . By (2.4), we see that y^* has infinitely many zeros, and $y(-\infty) = 0$, $|y^*(x)| = |u^*(-x)e^{\frac{x}{2}}| \leq \frac{1}{2}$, for $x \in (-\infty, 0]$. $\square$

Theorem 3 *For any integer $n \geq 0$, there is a unique solution $y^{(n)}$ to the following problem*

$$y'' - y' + y = y^3, \quad 0 < x < \infty,$$

$$y(0) = 0, \quad y(\infty) = 1,$$

y has precisely n zeros in $(0, \infty)$.

Proof. By Theorem 2, it is easy to check that

$$y^{(n)}(x) = y^*(x + x_n)$$

is a solution to this problem, where x_n is given in Theorem 2.

Now suppose there is another solution $\bar{y}^{(n)}$ to this problem. Let $0 = \bar{x}_0 < \bar{x}_1 < \bar{x}_2 < \dots < \bar{x}_n$ be the zeros of $\bar{y}^{(n)}$ in $[0, \infty)$. Then $y(s) = \bar{y}^{(n)}(s + \bar{x}_n)$, satisfies (2.8), (2.9), (2.10) with $x = s$. By Theorem 2, we have $y(s) = y^*(s)$, $-\infty < s < \infty$, or $\bar{y}^{(n)}(x) = y^*(x - \bar{x}_n)$. Then $\bar{x}_k - \bar{x}_n (k = 0, 1, \dots, n)$ are the zeros of y^* , $y^*(\bar{x}_k - \bar{x}_n) = \bar{y}^{(n)}(\bar{x}_k) = 0$, satisfying $-\bar{x}_n = \bar{x}_0 - \bar{x}_n < \bar{x}_1 - \bar{x}_n < \dots < \bar{x}_k - \bar{x}_n < \dots < \bar{x}_n - \bar{x}_n = 0$. Because $0, \bar{x}_k (k = 1, \dots, n)$ are the consecutive zeros of $\bar{y}^{(n)}$, $\bar{x}_k - \bar{x}_n (k = 0, 1, \dots, n)$ are the consecutive zeros of y^* . So we see that $x_n = -\bar{x}_n$. Thus $\bar{y}^{(n)}(x) = y^*(x - \bar{x}_n) = y^*(x + x_n) = y^{(n)}(x)$. So the theorem is proved. $\square$

3 Numerical Results

In this section, we present numerical computation results for some important values related to y^* . First of all, as seen in the previous sections, the most important number is a^* , which defines y^* . In the next section, we investigate the asymptotics of $y^*(x)$ as x approaches to $-\infty$ and as x approaches to ∞ . The following values will be used to represent the asymptotics,

$$b^* = \int_{-\infty}^0 e^{-\frac{s}{2}} \cos \frac{\sqrt{3}}{2} s (y^*(s))^3 ds, \quad (3.1)$$

$$c^* = \int_{-\infty}^0 e^{-\frac{s}{2}} \sin \frac{\sqrt{3}}{2} s (y^*(s))^3 ds, \quad (3.2)$$

$$d^* = \int_0^{\infty} e^s \left((1 - y^*(s))^2 - \frac{1}{3} (1 - y^*(s))^3 \right) ds. \quad (3.3)$$

(1) a^*

In Theorem 2 of [2], we obtained that

$$\left(\frac{1}{2} + 2 \sum_{n=1}^{N_1} \frac{b_n}{n+1} \right)^{\frac{1}{2}} < a^* < - \sum_{n=1}^{N_2} b_n,$$

for any $N_1 \geq 2$, and $N_2 \geq 1$. If we choose large numbers $N_1 = 170$, $N_2 = 170$, the *Mathematica* shows the following result

$$0.16871221576 \dots < a^* < 0.16871221594 \dots. \quad (3.4)$$

(2) $y^*(x)$ and $P(z)$

In last section, we have proved that $y^*(x)$ is the unique solution to problem (P). And it was also shown that $y^*(x)$ is the solution of the following initial value problem

$$y'' - y' + y = y^3, \quad -\infty < x < \infty, \quad (3.5)$$

$$y(0) = 0, \quad y'(0) = a^*. \quad (3.6)$$

So we can approach the function $y^*(x)$ by numerically solving the initial value problem with properly approximate value of a^* given above. The *Mathematica* gives the graph of $y^*(x)$. See Figure 1.

In section 4 and 5, we discussed the function $P(z)$ which is the solution to equation $PP' - P = z(z-1)(z-2)$, $z \in [0, 1]$ with $P(z) \sim -z$ as $z \rightarrow 0$.




Figure 1: Graph of $y^*(x)$ (solution of (P)). The value of $a^* = y^{*'}(0)$ is chosen $0.168712 \dots$.

As mentioned in [3], $P(z)$ gives the relation between $y^{*'}(x)$ and $y^*(x)$ for $x > 0$, which is

$$y^{*'}(x) = -P(1 - y^*(x)).$$

The graph of $P(z)$ is computed also by *Mathematica*, and given in Figure 2. The numerical computation for $P(z)$ shows that $P(1) \approx -0.1687\dots$, which matches the result (3.4).

(3) b^* , c^* , and d^*

The values of b^* , c^* , d^* will be used to represent the asymptotics of $y^*(x)$ as $x \rightarrow \pm\infty$, and the asymptotics of $f(r)$ as $r \rightarrow 0$ and $r \rightarrow \infty$ in the next section. We do not have further analytic properties for these three numbers since we have not found the analytic representation of y^* . Computer shows the approximate values of these three numbers based on




Figure 2: Graph of $P(z)$ as solution to the equation $PP' - P = z(z-1)(z-2)$, $z \in [0, 1]$ with $P(z_0) = -z_0$, where $z_0 = 10^{-40}$.

the numerical computation for $y^*(x)$,

$$b^* \approx \int_{-30}^0 e^{-\frac{s}{2}} \cos \frac{\sqrt{3}}{2} s (y^*(s))^3 ds \approx -0.0005497 \dots, \quad (3.7)$$

$$c^* \approx \int_{-30}^0 e^{-\frac{s}{2}} \sin \frac{\sqrt{3}}{2} s (y^*(s))^3 ds \approx 0.001939 \dots, \quad (3.8)$$

$$d^* \approx \int_0^{15} e^s \left((1 - y^*(s))^2 - \frac{1}{3}(1 - y^*(s))^3 \right) ds \approx 4.1728 \dots. \quad (3.9)$$

4 Asymptotics and Connection Formulas

We have seen in Sect. 2 that the problem (P) has a unique solution $y^*(x)$. Now, Let us consider the asymptotics of $y^*(x)$ as $x \rightarrow -\infty$, and $x \rightarrow \infty$.

Theorem 4 *The function $y^*(x)$ has the following asymptotics*

(i) *As $x \rightarrow -\infty$,*

$$y^*(x) = Ae^{\frac{x}{2}} \sin\left(\frac{\sqrt{3}}{2}x + \phi\right) + O(e^{\frac{3x}{2}}), \quad (4.1)$$

with the connection formula

$$A = \frac{2}{\sqrt{3}} \sqrt{(a^* - b^*)^2 + c^{*2}} \approx 0.196, \quad (4.2)$$

$$\phi = \arctan \frac{c^*}{a^* - b^*} \approx 0.0115 \approx 0.00375\pi. \quad (4.3)$$

(ii) *As $x \rightarrow 0$, there is*

$$y^*(x) \sim a^*x. \quad (4.4)$$

(iii) *As $x \rightarrow \infty$,*

$$y^*(x) = 1 - Be^{-x} + O(e^{-2x}), \quad (4.5)$$

with the connection formula

$$B = \frac{2 + a^*}{3} + d^* \approx 4.90. \quad (4.6)$$

Proof. Let us first consider (4.1). Recall that if we let

$$t = -x, u^*(t) = e^{-\frac{x}{2}} y^*(x),$$

then u^* satisfies for $t > 0$

$$u^{*''} + \frac{3}{4}u^* = e^{-t}u^{*3}(t),$$

and

$$\begin{aligned} u^*(t) &= -\frac{2}{\sqrt{3}}a^* \sin \frac{\sqrt{3}}{2}t + \frac{2}{\sqrt{3}} \int_0^t e^{-s} \sin \frac{\sqrt{3}}{2}(t-s) u^{*3}(s) ds \\ &= -\frac{2}{\sqrt{3}}a^* \sin \frac{\sqrt{3}}{2}t + \frac{2}{\sqrt{3}} \int_0^\infty e^{-s} \sin \frac{\sqrt{3}}{2}(t-s) u^{*3}(s) ds + O(e^{-t}), \end{aligned}$$

as $t \rightarrow \infty$. Direct calculation shows that

$$\begin{aligned} y^*(x) &= \frac{2}{\sqrt{3}}(a^* - b^*)e^{\frac{x}{2}} \sin \frac{\sqrt{3}}{2}x + \frac{2}{\sqrt{3}}c^*e^{\frac{x}{2}} \cos \frac{\sqrt{3}}{2}x + O(e^{\frac{3x}{2}}) \\ &= Ae^{\frac{x}{2}} \sin\left(\frac{\sqrt{3}}{2}x + \phi\right) + O(e^{\frac{3x}{2}}), \end{aligned}$$

where A and ϕ are given by (4.2),(4.3).

The asymptotics (4.4) is obvious. Finally let us prove (4.5). Let

$$y^*(x) = 1 - z(x).$$

It's easy to check that $z(x)$ satisfies the differential equation

$$z'' - z' - 2z = -3z^2 + z^3, \quad (4.7)$$

and integral equation

$$\begin{aligned} z(x) &= \frac{1 - a^*}{3} e^{2x} + \frac{2 + a^*}{3} e^{-x} + \\ &\quad \frac{e^{2x}}{3} \int_0^x e^{-2s} (-3z^2(s) + z^3(s)) ds - \frac{e^{-x}}{3} \int_0^x e^s (-3z^2(s) + z^3(s)) ds, \end{aligned}$$

since $z(0) = 1, z'(0) = -a^*$. Using equation (4.7) and Lemma 3.1 (iii) in [2], we have

$$\begin{aligned} &\int_0^\infty e^{-2s} (-3z^2(s) + z^3(s)) ds \\ &= \int_0^\infty e^{-2s} (z''(s) - z'(s) - 2z(s)) ds \\ &= -z'(0) - z(0) + ((-2)^2 + (-2) - 2) \int_0^\infty e^{-2s} z(s) ds \\ &= a^* - 1. \end{aligned}$$

Thus

$$\begin{aligned} z(x) &= \frac{2 + a^*}{3} e^{-x} - \frac{e^{-x}}{3} \int_0^x e^s (-3z^2(s) + z^3(s)) ds + O(e^{-2x}) \\ &= \left(\frac{2 + a^*}{3} + d^* \right) e^{-x} + O(e^{-2x}). \end{aligned}$$

So (4.5) holds. $\square$

Theorem 5 *Any solution $y(x)$ to the problem*

$$y'' - y' + y = y^3, \quad -\infty < x < \infty, \quad (4.8)$$

$$y(x) \rightarrow 0, \quad \text{as } x \rightarrow -\infty \quad (4.9)$$

$$y(x) \rightarrow 1, \quad \text{as } x \rightarrow \infty \quad (4.10)$$

has the representation

$$y(x) = y^*(x - \tau), \quad (4.11)$$

where τ is the largest zero point of $y(x)$. Hence the solution to (4.8), (4.9),(4.10) is unique up to a transition of x .

Proof. Let $y_1 = y, y_2 = y'$, and change equation (4.8) into the system

$$\begin{aligned} y_1' &= y_2, \\ y_2' &= y_2 - y_1 + y_1^3. \end{aligned}$$

It is not hard to see that the point $(0, 0)$ in the phase plane is a stable focus of the system (see [1]). If $y(x)$ is a solution to (4.8), (4.9) and (4.10), then $y(x)$ has infinitely many zeros. Because $y(\infty) = 1$, the zeros of $y(x)$ have upper bound, and the largest zero τ exists(finite). Let

$$y_1(x) = y(x + \tau).$$

Then $y_1(x)$ solves (P). By Theorem 2, $y_1(x) = y^*(x)$, and then (4.11) holds. $\square$

The solution to problem (P) is clear now. Let us come back to the original equation (1.1).

Theorem 6 *Any solution to (1.1), (1.2) and (1.3) has the representation*

$$f(r) = y^*(\log \frac{r}{r_0}), \quad (4.12)$$

where r_0 is the largest zero of $f(r)$. With a^*, b^*, c^* and d^* defined by (1.4), (3.1), (3.2) and (3.3) respectively, $f(r)$ has the following asymptotics.

(i) As $r \rightarrow 0$,

$$f(r) = A \left(\frac{r}{r_0} \right)^{\frac{1}{2}} \sin \left(\frac{\sqrt{3}}{2} \log \frac{r}{r_0} + \phi \right) + O(r^{\frac{3}{2}}), \quad (4.13)$$

where again

$$A = \frac{2}{\sqrt{3}} \sqrt{(a^* - b^*)^2 + c^{*2}} \approx 0.196, \quad (4.14)$$

$$\phi = \arctan \frac{c^*}{a^* - b^*} \approx 0.0115 \approx 0.00375\pi. \quad (4.15)$$

(ii) As $r \rightarrow r_0$,

$$f(r) \sim a^* \log \frac{r}{r_0}, \quad (4.16)$$

(iii) As $r \rightarrow \infty$,

$$f(r) = 1 - \frac{Br_0}{r} + O\left(\frac{1}{r^2}\right), \quad (4.17)$$

where

$$B = \frac{2 + a^*}{3} + d^* \approx 4.90. \quad (4.18)$$

Proof. By Theorem 4 and Theorem 5, using $r = e^x, r_0 = e^\tau, \log \frac{r}{r_0} = x - \tau, f(r) = y(x)$, direct calculations show the results. $\square$

References

- [1] Perko, L.: *Differential Equations and Dynamical Systems*, 2nd ed. Springer-Verlag, New York, 1996.
- [2] Wang, C. B.: *Boundary value problem for $r^2 d^2 f/dr^2 + f = f^3$ (I): existence and uniqueness*, preprint.
- [3] Wang, C. B.: *Boundary Value Problem for $r^2 d^2 f/dr^2 + f = f^3$ (II): connection formula*, preprint.